



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

he and Descartes carried on an intimate correspondence, each being a great admirer of the other. Little, however, could either have guessed that Christian, then only a child of three, would one day far outrank his father and would, in point of mathematical ability, rival his father's distinguished friend.

QUESTIONS AND DISCUSSIONS.

EDITED BY W. A. HURWITZ, Cornell University, Ithaca, N. Y.

DISCUSSIONS.

To a question raised by the editor in connection with Professor Light's "Note on curves whose evolutes are similar curves" [1920, 303], as to the existence of other curves possessing the stated property, an answer is given below by Mr. Franklin, who derives in a simple way the result, stated by Puiseux, that an infinite set of such curves exists. A rather interesting incidental feature is found at the end of the paper, in the construction of a sort of geometrical "sum" of two curves as the locus of the sum (as determined by vectors starting at the origin) of points on the two curves at which tangents are parallel. The especial interest lies in the fact that exactly this process of composition of curves has been mentioned recently in an entirely different connection by Professor W. B. Carver¹; the curves to which he applies the process are algebraic, while those of Mr. Franklin are transcendental.

Proofs of the law of tangents in plane trigonometry have been given recently by Cheney [1920, 53], Lovitt [1920, 465] and Epperson [1921, 71]. In a letter to the editor, from which an extract is printed, Professor Mathews calls attention to various other proofs of the law. It is clear, as Professor Mathews states, that other new proofs can be devised in considerable number from a suitable figure; it therefore seems desirable to bar consideration of further proofs, unless they involve new principles.

As the third discussion, we print a note on the nature of expository papers for presentation to the Association.

I. ON CURVES WHOSE EVOLUTES ARE SIMILAR CURVES.

BY PHILIP FRANKLIN, Princeton University.

In the July-September number of the MONTHLY [1920, 303] there appeared a discussion of curves whose evolutes are similar to themselves, by Professor Light. He found that the only curves having this property and having their intrinsic equations of the special type $AR^n + BS^m + C = 0$ were the logarithmic spiral and the cycloidal curves. The editor inquired whether any other curves possessed this property.

¹ "The failure of the Clifford chain," *American Journal of Mathematics*, vol. 42 (1920), p. 167.